



AMH gene

anti-Müllerian hormone

Normal Function

The *AMH* gene provides instructions for making a protein that is involved in male sex differentiation. During development of male fetuses, the AMH protein is produced and released (secreted) by cells of the testes. The secreted protein attaches (binds) to its receptor, which is found on the surface of Müllerian duct cells. The Müllerian duct, found in both male and female fetuses, is the precursor to the female reproductive organs. Binding of the AMH protein to its receptor induces self-destruction (apoptosis) of the Müllerian duct cells. As a result, the Müllerian duct breaks down (regresses) in males. In females, who do not produce the AMH protein during fetal development, the Müllerian duct becomes the uterus and fallopian tubes.

Health Conditions Related to Genetic Changes

persistent Müllerian duct syndrome


Persistent Müllerian duct syndrome type 1, a disorder of sexual development that affects males, is caused by mutations in the *AMH* gene. Males with this condition have female reproductive organs in addition to normal male reproductive organs. At least 38 mutations in the *AMH* gene have been identified in people with persistent Müllerian duct syndrome type 1. Most mutations change single protein building blocks (amino acids) in the AMH protein. Other mutations result in a premature stop signal that leads to an abnormally short protein. Still other mutations delete regions of DNA from the *AMH* gene, which changes the instructions for the protein.

The mutated AMH protein cannot be released from the cells of the testes or cannot bind to the receptor on the Müllerian duct cells. As a result, the Müllerian duct cells never receive the signal for apoptosis. The Müllerian duct persists and becomes a uterus and fallopian tubes. Because the AMH protein is not involved in the formation of male reproductive organs, affected males also have male reproductive organs.

Chromosomal Location

Cytogenetic Location: 19p13.3, which is the short (p) arm of chromosome 19 at position 13.3

Molecular Location: base pairs 2,249,114 to 2,252,073 on chromosome 19 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- anti-Muellerian hormone
- MIF
- MIS
- muellerian-inhibiting factor
- muellerian-inhibiting substance
- Mullerian inhibiting factor
- Mullerian inhibiting substance

Additional Information & Resources

Educational Resources

- An Introduction to Genetic Analysis (seventh edition, 2000): Development of the Urogenital System
<https://www.ncbi.nlm.nih.gov/books/NBK21846/figure/A3724/>
- StemBook (2008): Müllerian Duct Formation and Neonatal Development
<https://www.ncbi.nlm.nih.gov/books/NBK27042/#uterinestemcells.sec2-1>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28AMH%5BTIAB%5D%29+OR+%28anti-Mullerian+hormone%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

OMIM

- ANTI-MULLERIAN HORMONE
<http://omim.org/entry/600957>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_AMH.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=AMH%5Bgene%5D>
- HGNC Gene Family: Endogenous ligands
<http://www.genenames.org/cgi-bin/genefamilies/set/542>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=464
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/268>
- UniProt
<http://www.uniprot.org/uniprot/P03971>

Sources for This Summary

- OMIM: ANTI-MULLERIAN HORMONE
<http://omim.org/entry/600957>
- Belville C, Van Vlijmen H, Ehrenfels C, Pepinsky B, Rezaie AR, Picard JY, Josso N, di Clemente N, Cate RL. Mutations of the anti-mullerian hormone gene in patients with persistent mullerian duct syndrome: biosynthesis, secretion, and processing of the abnormal proteins and analysis using a three-dimensional model. *Mol Endocrinol*. 2004 Mar;18(3):708-21. Epub 2003 Dec 12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14673134>
- Josso N, Belville C, di Clemente N, Picard JY. AMH and AMH receptor defects in persistent Müllerian duct syndrome. *Hum Reprod Update*. 2005 Jul-Aug;11(4):351-6. Epub 2005 May 5. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15878900>

- Josso N, Picard JY, Imbeaud S, di Clemente N, Rey R. Clinical aspects and molecular genetics of the persistent müllerian duct syndrome. Clin Endocrinol (Oxf). 1997 Aug;47(2):137-44. Review. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/9302384>
 - Rey R. Anti-Müllerian hormone in disorders of sex determination and differentiation. Arq Bras Endocrinol Metabol. 2005 Feb;49(1):26-36. Epub 2006 Mar 16. Review. *Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16544032>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/AMH>

Reviewed: March 2011

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services